Arthritis - (Calculated Variable)*
Kentucky BRFSS 2000

Geographic Groups	Total Respond.#	Yes			No		
		N	%	C.I.(95%)	N	%	C.I.(95%)
TOTAL	6326	2837	40.0	(38.3-41.7)	3489	60.0	(58.3-61.7)
ADDs							
Barren River	436	189	40.0	(34.5-45.6)	247	60.0	(54.4-65.5)
Big Sandy	340	178	48.7	(42.7-54.7)	162	51.3	(45.3-57.3)
Bluegrass	500	207	39.6	(34.9-44.5)	293	60.4	(55.5-65.1)
Buffalo Trace	330	150	40.6	(34.8-46.6)	180	59.4	(53.4-65.2)
Cumberland Valley	395	202	45.8	(40.2-51.6)	193	54.2	(48.4-59.8)
FIVCO	460	226	46.5	(40.3-52.8)	234	53.5	(47.2-59.7)
Gateway	447	207	41.6	(36.4-47.1)	240	58.4	(52.9-63.6)
Green River	377	158	37.6	(32.3-43.1)	219	62.4	(56.9-67.7)
Kentucky River	421	225	47.7	(42.2-53.3)	196	52.3	(46.7-57.8)
KIPDA	465	189	38.4	(33.5-43.6)	276	61.6	(56.4-66.5)
Lake Cumberland	405	188	41.3	(35.8-47.0)	217	58.7	(53.0-64.2)
Lincoln Trail	430	174	36.9	(32.0-42.0)	256	63.1	(58.0-68.0)
Northern Kentucky	432	164	36.3	(31.4-41.4)	268	63.7	(58.6-68.6)
Pennyrile	425	185	39.2	(34.2-44.5)	240	60.8	(55.5-65.8)
Purchase	463	195	37.1	(32.4-42.1)	268	62.9	(57.9-67.6)

*For this calculated variable 'Yes' means that respondents have eitherchronic joint symptoms or have ever been told by a doctor that they have arthritis. Response is generalized to the state population

#Use caution in interpreting data for total respondents less than 50. N = Cell Size, NH = Non-Hispanic, % = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level).

Percentages are weighted to population characteristics.

Unless stated differently above, Missing, 'Don't know', and Refused responses are excluded.